

## Lyme Disease Association Analysis of CDC 2007 Reported Lyme Disease Cases

State	1993	2000	2006	2007	% Increase from 2006 to 2007	% of total 2007 US reported cases	States ranked by 2007 reported cases (*tie)	% Increase from 2000 to 2007	% Increase from 1993 to 2007
Alabama	4	6	11	13	18%	0.05%	28	117%	225%
Alaska	0	2	3	10	233%	0.04%	*31	400%	1000%
Arizona	0	2	10	2	-80%	0.01%	*39	0%	300%
Arkansas	8	7	0	1	800%	0.00%	*40	-86%	0%
California	134	96	85	75	-12%	0.27%	20	-22%	-44%
Colorado	0	0	0	0	0%	0.00%	N/A	0%	0%
Connecticut	1350	3773	1788	3058	71%	11.14%	4	-19%	127%
DC	2	11	62	116	87%	0.42%	17	955%	5700%
Delaware	143	167	482	715	48%	2.61%	11	328%	400%
Florida	30	54	34	30	-12%	0.11%	26	-44%	0%
Georgia	44	0	8	11	38%	0.04%	30	1100%	-75%
Hawaii	1	0	0	Not Notifiable	NA	NA	NA	NA	NA
Idaho	2	4	7	9	29%	0.03%	32	125%	350%
Illinois	19	35	110	149	35%	0.54%	14	326%	684%
Indiana	32	23	26	55	112%	0.20%	21	139%	72%
Iowa	8	34	97	123	27%	0.45%	16	262%	1438%
Kansas	54	17	4	8	100%	0.03%	33	-53%	-85%
Kentucky	16	13	7	6	-14%	0.02%	*35	-54%	-63%
Louisiana	3	8	1	2	100%	0.01%	*39	-75%	-33%
Maine	18	71	338	529	57%	1.93%	12	645%	2839%
Maryland	180	688	1248	2576	106%	9.39%	6	274%	1331%
Massachusetts	148	1158	1432	2988	109%	10.89%	5	158%	1919%
Michigan	23	23	55	51	-7%	0.19%	23	122%	122%
Minnesota	141	465	914	1238	35%	4.51%	8	166%	778%
Mississippi	0	3	3	1	-67%	0.00%	*40	-67%	200%
Missouri	108	47	5	10	100%	0.04%	*31	-79%	-91%
Montana	0	0	1	4	300%	0.01%	37	700%	700%
Nebraska	6	5	11	7	-36%	0.03%	*34	40%	17%
Nevada	5	4	4	15	275%	0.05%	27	275%	200%
New Hampshire	15	84	617	896	45%	3.26%	10	967%	5873%
New Jersey	786	2459	2432	3134	29%	11.42%	3	27%	299%
New Mexico	2	0	3	5	67%	0.02%	36	500%	150%
New York	2818	4329	4460	4165	-7%	15.18%	1	-4%	48%
North Carolina	86	47	31	53	71%	0.19%	22	13%	-38%
North Dakota	2	2	7	12	71%	0.04%	*29	500%	500%
Ohio	30	61	43	33	-23%	0.12%	24	-46%	10%
Oklahoma	19	1	0	1	100%	0.00%	*40	0%	-95%
Oregon	8	13	7	6	-14%	0.02%	*35	-54%	-25%
Pennsylvania	1085	2343	3242	3994	23%	14.55%	2	70%	268%
Rhode Island	272	675	308	177	-43%	0.64%	13	-74%	-35%
South Carolina	9	25	20	31	55%	0.11%	*25	24%	244%
South Dakota	0	0	1	0	-100%	0.00%	N/A	0%	0%
Tennessee	20	28	15	31	107%	0.11%	*25	11%	55%
Texas	48	77	29	87	200%	0.32%	18	13%	81%
Utah	2	3	5	7	40%	0.03%	*34	133%	250%
Vermont	12	40	105	138	31%	0.50%	15	245%	1050%
Virginia	95	149	357	959	169%	3.49%	9	544%	909%
Washington	9	9	8	12	50%	0.04%	*29	33%	33%
West Virginia	50	35	28	84	200%	0.31%	19	140%	68%
Wisconsin	401	631	1466	1814	24%	6.61%	7	187%	352%
Wyoming	9	3	1	3	200%	0.01%	38	0%	-67%
<b>U.S. TOTAL</b>	<b>8,257</b>	<b>17,730</b>	<b>19,931</b>	<b>27,444</b>	<b>38%</b>	<b>N/A</b>	<b>N/A</b>	<b>55%</b>	<b>232%</b>

Note: Updated Numbers reported by CDC on 9/19/08. CDC says Lyme disease is under reported and that only about 10% of cases that meet CDC surveillance criteria are actually reported to CDC. (E.g. New York 4,165 reported cases = 41,650 probable cases meeting CDC criteria) Source Data compiled: CDC pub data (MMWR) ©2008 Lyme Disease Association LDA PO Box 1438 Jackson, NJ 08527  
[www.LymeDiseaseAssociation.org](http://www.LymeDiseaseAssociation.org) 888 366 6611